

Solving Quadratic Equations Algebraically

Definition of a Quadratic Equation

A *quadratic*, or *second degree*, equation is one that can be written in the form

$$ax^2 + bx + c = 0$$

for real constants a , b , and c , with $a \neq 0$.

NOTE This chapter considers only real solutions, that is, solutions that are real numbers.

Techniques Used to Solve Quadratic Equations

There are four techniques normally used to algebraically find exact solutions of quadratic equations. Techniques that can be used to solve *some* quadratic equations include

- factoring $x^2 - 4 = 0$ $(x-2)(x+2) = 0$ $x = 2, -2$
- taking the square root of both sides of an equation $\sqrt{x^2} = \sqrt{9}$ $x = \pm 3$

2.2

Techniques that can be used to solve *all* quadratic equations include

2.1

- completing the square
- using the quadratic formula
- using graphing calc.

Solving Quadratic Equations by Factoring

The factoring method of solving quadratic equations is based on the Zero Product Property of real numbers.

$$(x+5)(x-3) = 0$$

If a product of real numbers is zero, then at least one of the factors is zero. In other words,

If $ab = 0$, then $a = 0$ or $b = 0$ (or both).

1-73 e.o.o.

The Zero Product Property

Example 1 Solving a Quadratic Equation by Factoring

Solve $3x^2 - x = 10$ by factoring.

$$y_1 = 3x^2 - x - 10$$

$$3x^2 - x = 10$$

$$3x^2 - x - 10 = 0$$

$$(3x + 5)(x - 2) = 0$$

$$-6x + 5x - 1x$$

$$3x + 5 = 0$$

$$x = 2$$

$$x = -5/3$$

These are like 1-12.




Figure 2.2-1

These are like 13-24.

Taking the Square Root of Both Sides of an Equation

Example 2 Solving $ax^2 = b$

Solve $\frac{3x^2}{3} = \frac{9}{3}$

$$\sqrt{x^2} = \sqrt{3}$$

$$x = \pm\sqrt{3}$$

$$4x^2 - 1 = 23$$

$$\frac{4x^2}{4} = \frac{24}{4}$$

$$\sqrt{x^2} = \sqrt{6}$$

$$x = \pm\sqrt{6}$$

CAUTION

When taking the square root of both sides of an equation, remember to write \pm on one side of the equation.

$$\sqrt{x^2} = \sqrt{12}$$

$$x = \pm 2\sqrt{3}$$

The method of taking the square root of both sides of an equation can be used to solve equations of the form $a(x - h)^2 = k$.

Example 3 Solving $a(x - h)^2 = k$

Solve $\frac{2(x+4)^2}{2} = \frac{6}{2}$

$$\sqrt{(x+4)^2} = \sqrt{3}$$

$$x+4 = \pm\sqrt{3}$$

$$x = -4 \pm \sqrt{3}$$

$$2(x-1)^2 + 8 = 30$$

$$\frac{2(x-1)^2}{2} = \frac{22}{2}$$

$$\sqrt{(x-1)^2} = \sqrt{11}$$

$$x-1 = \pm\sqrt{11}$$

$$x = 1 \pm \sqrt{11}$$

Example 4 Solving a Quadratic Equation by Completing the Square

Solve $2x^2 - 8x + 4 = 0$ by completing the square.

$$\frac{2x^2}{2} - \frac{8x}{2} + \frac{4}{2} = \frac{0}{2}$$

$$x^2 - 4x + 2 = 0$$

$$x^2 - 4x + 4 = -2 + 4$$

$$\sqrt{(x-2)^2} = \sqrt{2}$$

$$x-2 = \pm\sqrt{2}$$

$$x = 2 \pm \sqrt{2}$$

$$-\frac{4}{2} = (-2)^2 = 4$$

Like 25-52

$$x^2 - 12x - 6 = 0$$

$$x^2 - 12x + 36 = 6 + 36$$

$$\sqrt{(x-6)^2} = \sqrt{42}$$

$$x-6 = \pm\sqrt{42}$$

$$x = 6 \pm \sqrt{42}$$

The Quadratic Formula

The solutions of the quadratic equation $ax^2 + bx + c = 0$ are

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Because the quadratic formula can be used to solve *any* quadratic equation, it should be memorized.

Solving Quadratic Equations Algebraically

B.M. 1-73 e00

Definition of a Quadratic Equation

A *quadratic*, or *second degree*, equation is one that can be written in the form

$$ax^2 + bx + c = 0$$

for real constants a , b , and c , with $a \neq 0$.

NOTE This chapter considers only real solutions, that is, solutions that are real numbers.

Techniques Used to Solve Quadratic Equations

There are four techniques normally used to algebraically find exact solutions of quadratic equations. Techniques that can be used to solve *some* quadratic equations include

- factoring
- taking the square root of both sides of an equation

Techniques that can be used to solve *all* quadratic equations include

- completing the square
- using the quadratic formula

$$(5x+4)(x-6)=0$$

\downarrow \downarrow
 $-4/5$ 6

Solving Quadratic Equations by Factoring

The factoring method of solving quadratic equations is based on the Zero Product Property of real numbers.

$$(x-3)(x+2)=0$$

If a product of real numbers is zero, then at least one of the factors is zero. In other words,

If $ab = 0$, then $a = 0$ or $b = 0$ (or both).

$$x = 3 \qquad x = -2$$

The Zero Product Property

Example 1 Solving a Quadratic Equation by Factoring

Solve $3x^2 - x = 10$ by factoring. get a 0 on one side

These are like 1-12.

$$3x^2 - x - 10 = 0$$

$$(3x + 5)(x - 2) = 0$$

$$(3x + 5)(x - 2) = 0$$

$$3x + 5 = 0$$

$$3x = -5$$

$$x = -\frac{5}{3}$$

$$x - 2 = 0$$

$$x = 2$$




Figure 2.2-1

The Quadratic Formula

The solutions of the quadratic equation $ax^2 + bx + c = 0$ are

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Because the quadratic formula can be used to solve *any* quadratic equation, it should be memorized.

Example 5 Solving a Quadratic Equation by Using the Quadratic Formula

Like 29-40

Solve $x^2 + 3 = -8x$ by using the quadratic formula.

$$x^2 + 8x + 3 = 0$$

$$a=1 \quad b=8 \quad c=3$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-8 \pm \sqrt{64 - 4(1)(3)}}{2}$$

$$x = \frac{-8 \pm \sqrt{64 - 12}}{2}$$

$$x = \frac{-8 \pm \sqrt{52}}{2}$$

$$x = \frac{-8 \pm 2\sqrt{13}}{2}$$

$$\Rightarrow -4 \pm \sqrt{13}$$

Best answer

Handwritten calculation for $\sqrt{52}$:

$$\begin{array}{r} 52 \\ 2 \overline{) 52} \\ \underline{40} \\ 120 \\ \underline{104} \\ 160 \\ \underline{156} \\ 40 \end{array}$$

The Discriminant

The expression $b^2 - 4ac$ in the quadratic formula, called the discriminant, can be used to determine the *number* of real solutions of the equation $ax^2 + bx + c = 0$.

Real Solutions of a Quadratic Equation

Discriminant Value	Number of Real Solutions of $ax^2 + bx + c = 0$
$b^2 - 4ac > 0$ <i>pos.</i>	2 distinct real solutions
$b^2 - 4ac = 0$	1 distinct real solution
$b^2 - 4ac < 0$ <i>negative</i>	0 real solutions

The discriminant can be used to determine if an equation has no real solutions without completing all computations.

Example 6 Determining the Number of Solutions by Using the Discriminant

Like 41-46

Solve $2x^2 = -x - 3$.

$$2x^2 + 1x + 3 = 0$$

\uparrow \uparrow \uparrow
 a b c

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$d = \begin{cases} b^2 - 4ac \\ 1^2 - 4(2)(3) \\ 1 - 24 = -23 \end{cases}$$

How many sol.?
0

Polynomial Equations in Quadratic Form

Like 61-68

Example 7 Solving an Equation in Quadratic Form

Solve 4.④ $-13x^2 + 3 = 0$.

$$4x^2 - 13x + 3 = 0$$

$$(4x^2 - 1)(x^2 - 3) = 0$$

$$\begin{aligned} 4x^2 - 1 &= 0 \\ \sqrt{x^2} &= \sqrt{\frac{1}{4}} \\ x &= \pm \frac{1}{2} \end{aligned}$$

$$\begin{aligned} x^2 - 3 &= 0 \\ \sqrt{x^2} &= \sqrt{3} \\ x &= \pm \sqrt{3} \end{aligned}$$

~~$$(4x - 3)(x - 1)$$

$$4x^2 - 4x - 3x + 3$$

$$-7x$$~~

1-73 eoo

In Exercises 69-72, find a number k such that the given equation has exactly one real solution.

Like 71, 73

69. $x^2 + kx + 25 = 0$

$$b^2 - 4ac = 0$$

$$k^2 - 4(1)(25) = 0$$

$$k^2 - 100 = 0 \Rightarrow \sqrt{k^2} = \sqrt{100}$$

$$k = \pm 10$$

$$(k - 10)(k + 10) = 0$$

$$k = 10 \quad k = -10$$

Polynomial Equations in Quadratic Form

Like 61-68

Example 7 Solving an Equation in Quadratic Form

Solve $4x^4 - 13x^2 + 3 = 0$.

$$4x^4 - 13x^2 + 3 = 0$$

$$(4x^2 - 1)(x^2 - 3) = 0$$

$$\downarrow$$
$$4x^2 - 1 = 0$$

$$\sqrt{x^2} = \sqrt{\frac{1}{4}}$$

$$x = \pm \frac{1}{2}$$

$$x^2 - 3 = 0$$

$$\sqrt{x^2} = \sqrt{3}$$

$$x = \pm \sqrt{3}$$

In Exercises 69–72, find a number k such that the given equation has exactly one real solution.

Like 71

69. $x^2 + kx + 25 = 0$

$$\downarrow b^2 - 4ac = 0$$

$k = b$ value

$$b^2 - 4ac = 0$$

$$k^2 - 4(1)(25) = 0$$

$$k^2 - 100 = 0$$

$$(k - 10)(k + 10) = 0$$

$$k = 10 \quad k = -10$$

$$\rightarrow \sqrt{k^2} = \sqrt{100}$$
$$k = \pm 10$$

